Published by:



## JOURNAL OF MICROBIOLOGY, BIOTECHNOLOGY AND FOOD SCIENCES

HOME ARTICLE IN PRESS CURRENT ISSUE NEW ARCHIVE OLD ARCHIVE CONTACT TOOLS ISSN: 1338-5178



## FOR AUTHORS

SUBMIT MANUSCRIPTS

INSTRUCTIONS FOR AUTHORS

EDITORIAL MEMBERS

INDEXING AND PARTNERSHIPS

**STATEMENTS** 

SEARCH ON WEB OF SCIENCE

IMPACT FACTOR

JOURNAL METRICS

AGROBIOTECH CENTRE



This work is licensed under a Creative Commons Attribution 4.0 International License

Current issue

## NUCLEOTIDE DEGRADATION PRODUCTS, TOTAL VOLATILE BASIC NITROGEN, SENSORY AND MICROBIOLOGICAL QUALITY OF NILE PERCH (LATES NILOTICUS) FILLETS UNDER CHILLED STORAGE

Back to full issue: October - November 2012, vol. 2, no. 2

pages: 653-666

Authors: Andrew Kiri Amegovu, Mohammed Luyima Sserunjogi, Patrick Ogwok, Vincent Makokha

Article type: Food Sciences of Food Sciences

Abstract: Degradation products of adenosine nucleotide and total volatile basic nitrogen (TVBN) concentration provide means of ascertaining freshness of commercial fish products. A complementary sensory analysis has also been adopted by export markets for assessing the quality of fresh fish. Nucleotide breakdown products and TVBN was determined in fresh fillets from beach seined and gill netted Nile perch, a highly commercialized freshwater fish from Lake Victoria (Uganda), under chilled storage. Microbiological and sensory qualities were also evaluated. Total plate and Pseudomonas spp. counts positively correlated with TVBN. Basing on sensory, microbiological and biochemical attributes of the fillets, shelf-life of gill netted Nile perch was lower (13 days) than that of the beach seined (17 days). Fillets of beach seined Nile perch have a better keeping quality than that of the gill netted.

Keywords: Nile perch, nucleotides, TVBN, microbiological, sensory, quality

XMLs: | NLM DTD xml | Copernicus xml |

Full text pdf download link: Issue navigation: October - November 2012, vol. 2, no. 2:



prev. article |p. 640-652|

next article |p. 667-683|